

**SINGULAR INNER FUNCTION AND
ANALYTIC BOUNDED POINT EVALUATION**

Kifah Al-Hami

Mathematics Department

College of Science

University of Bahrain, 32038, BAHRAIN

Abstract: In Section 1, I constructed a singular inner function which is an upper bound for a given outer function, i.e. replacing an outer function with a constant multiple of a singular inner

In Section 2, I proved how unit disk containment is not effected for the analytic bounded point evaluation by a restricted measure.

AMS Subject Classification: 53C20, 53B20

Key Words: outer function, singular inner function, Carleson set, bounded point evaluation

**1. Construction of a Singular Inner Function
from a Given Outer One**

Lets start first by introducing the following three definitions :

Definition 1.1. A function F defined and analytic in D is called an outer function if it has the form

$$F(z) = \chi \exp\left[\frac{1}{2\pi} \int_0^{2\pi} \frac{e^{it} + z}{e^{it} - z} \log k(t) dt\right] \quad (z \in D),$$

where k is a non-negative Lebesgue measurable function on ∂D , $\log(k(t)) \in L^1(m)$ and χ is a complex number of modulus 1

Definition 1.2. A function S_μ defined and analytic in D is called a singular inner function if it has the form

$$S_\mu(z) = \exp\left[-\int_{\partial D} \frac{\xi+z}{\xi-z} d\mu(z)\right] \quad (z \in D),$$

where μ is a finite non-negative Borel measure on ∂D that is singular with respect to m

Definition 1.3. A subset E of ∂D is called a Carleson set if:

- 1) E is closed;
- 2) E is of Lebesgue measure zero ; $|E| = 0$;
- 3) $\sum_{\nu} |I_\nu| \log \frac{1}{|I_\nu|} < \infty$, where $\{I_\nu\}$ are the complementary arcs of E

Now, the following Theorem which appears in [1] is of great importance for the Theorem that follows

Theorem 1.4. Let Ω be any Jordan domain in D , with $m(\partial D \cap \partial\Omega) = 0$, let $0 \leq h \in L^1(dm)$ and let $P_z(\zeta)$ be the Poisson kernel on ∂D evaluated at z in Ω . Then, for any $\epsilon > 0$, there exists a finite measure μ with support in ∂D such that $\mu \perp m$, and $\mu(E) = 0$ for every Carleson set E in ∂D and $|\int_{\partial D} (P_z(\zeta)d\mu - P_z(\zeta)hdm)| \leq \epsilon$ for all z in Ω .

In this Theorem we construct a special singular inner function which is an upper bound for a given outer function

Theorem 1.5. Let Ω be any Jordan domain in D , with $m(\partial D \cap \partial\Omega) = 0$ and let F be an outer function on D ; $F \neq 0$. Then there exists a singular inner function S_μ , with $\mu(E) = 0$ for every Carleson set E in ∂D , such that $|S_\mu(z)| \leq 3 \cdot |F(z)|$ for all z in Ω .

Proof. Since F is an outer function, $|F(z)| = e^{\int_{\partial D} P_z(\zeta) \log h(\zeta) dm(\zeta)}$ without loss of generality we can assume that $0 \leq h \leq 1$ on ∂D . Now, by Theorem 1.4, there exists a finite positive Borel measure μ such that $\mu \perp m$, $\mu(E) = 0$ for every Carleson set $E \subseteq \partial D$ and $|\int_{\partial D} (P_z(\zeta)d\mu(\zeta) - P_z(\zeta)) \log h(\zeta) dm(\zeta)| \leq 1$ for all z in Ω . Let S_μ be the corresponding singular inner function. Then $|S_\mu(z)| = e^{-\int_{\partial D} P_z(\zeta)d\mu(\zeta)}$ and therefore, $|S_\mu(z)/F(z)| = e^{-\int_{\partial D} (P_z(\zeta)d\mu - \int_{\partial D} P_z(\zeta) \log(h) dm)} \leq e^{\int_{\partial D} (P_z(\zeta)|\log(h)| dm - P_z(\zeta)d\mu)} \leq 3$ for all z in Ω . □

2. The set of Overconvergence $abpe(P^t(\mu))$

We now turn to the topic of point evaluations and establish some "overconvergence" results

With μ and $P^t(\mu)$ as before, a complex number z is called a bounded point evaluation for $P^t(\mu)$ if there is a constant M such that $|p(z)| \leq M \cdot \|p\|_{L^t(\mu)}$ for all polynomials p ; the collection of all such points is denoted $bpe(P^t(\mu))$. If $z \in C$ and there are positive constants M and r such that $|p(w)| \leq M \cdot \|p\|_{L^t(\mu)}$ whenever p is a polynomial and $|w - z| < r$, then we call z an analytic bounded point evaluation for $P^t(\mu)$; the set of all points z of this type is denoted by $abpe(P^t(\mu))$. Notice that $abpe(P^t(\mu))$ is an open subset of $bpe(P^t(\mu))$ and, by the Maximum Modulus Theorem, each component of $abpe(P^t(\mu))$ is simply connected. If $z \in abpe(P^t(\mu))$, then by the Hahn-Banach and Riesz Representation Theorems, there exists K , in $L^s(\mu)$ ($1/s + 1/t = 1$) such that $p(z) = \int p(\xi)K_z(\xi)d\mu(\xi)$ for each polynomial p . For f in $P^t(\mu)$, define \hat{f} on $bpe(P^t(\mu))$ by $\hat{f}(z) = \int f(\xi)K_z(\xi)d\mu(\xi)$. Observe that $\hat{f} = f.a.e.\mu$ on $bpe(P^t(\mu))$ and in fact $z \mapsto \hat{f}(z)$ is analytic on $abpe(P^t(\mu))$. The set $abpe(P^t(\mu))$ support (μ) can be thought of as a set of overconvergence for $P^t(\mu)$. We now establish two results that address this topic of overconvergence.

Lemma 2.1. *Let μ be a finite, positive Borel measure with compact support in C . Let K be a compact subset of $abpe(P^t(\mu))$ and let $M = \sup\{|p(z)|^t : z \in K\}$, $p \in P$ and $\|p\|_{L^t(\mu)}^t = 1$. If $\mu(K) < 1/M$, then $\|p\|_{L^t(\mu)} \leq 1/(1 - M \cdot \mu(K))^{1/t} \cdot \|P\|_{L^t(\mu_{(C/K)})}$ for every polynomial p .*

Proof. By the definition of M , $|p(z)|^t \leq M \cdot \|p\|_{L^t(\mu)}^t$ for every polynomial p . Therefore, $\int_C |p|^t d\mu = \int_{C/K} |p|^t d\mu + \int_K |p|^t d\mu \leq \int_{C/K} |p|^t d\mu + M \cdot \|p\|_{L^t(\mu)}^t \mu(K)$, and hence $(1 - M \cdot \mu(K)) \cdot \int_C |p|^t d\mu \leq \int_{C/K} |p|^t d\mu$. Consequently, $\int_C |p|^t d\mu \leq \frac{1}{(1 - M \cdot \mu(K))} \int_{C/K} |p|^t d\mu$ □

Proposition 2.2. *Let μ be a finite, positive Borel measure with compact support in C such that $D \subseteq abpe(P^t(\mu))$. If K is a compact subset of D , then $D \subseteq abpe(P^t(\mu_{(C/K)}))$*

Proof. Let $\rho = \max\{|z| : z \in K\}$. Now choose r such that $\rho \leq r \leq 1$. Since $\{z : |z| \leq r\}$ is a compact subset of $abpe(P^t(\mu))$, there exists $M_r > 1$ such that $|p(z)| \leq M_r \cdot \|p\|_{L^t(\mu)}$ whenever $|z| \leq r$ and $p \in P$. Let $f_n(z) = (\frac{z}{r})^n$. Now, if $|z| = r$ and p is a polynomial, then $|p(z)| = |p(z) \cdot f_n(z)| \leq M_r \|p f_n\|_{L^t(\mu)} = M_r \|p\|_{L^t(\mu_n)}$ (where $d\mu_n = |f_n|^t d\mu$). Since $f_n \rightarrow 0$ uniformly on K , there exists N such that $\mu_N(K) \cdot M_r \leq 1/2^t$. So, by Lemma 2.1, $|p(z)| \leq$

$2M_r \|p\|_{L^t(\mu_N(C/K))}$ whenever $p \in P$ and $|z| = r$. Now, there exists $c > 0$ such that $\mu_N \leq c \cdot \mu$ and therefore $|p(z)| \leq 2cM_r \cdot \|p\|_{L^t(\mu_N(C/K))}$ whenever $p \in P$ and $|z| = r$. By the Maximum Modulus Theorem, $\{z : |z| < r\} \subseteq \text{abpe}(P^t(\mu_{(C/K)}))$. Since $\rho < r < 1$ is arbitrary, we conclude that $D \subseteq \text{abpe}(P^t(\mu_{(C/K)}))$ \square

References

- [1] K. Al-Hami, *Weak-Star convergence of measure*, International Journal of pure and applied Math. Vol. 88, No. 4, (2013).