

**AUTOMORPHISMS OF THE UNIT GROUPS OF SQUARE
RADICAL ZERO FINITE COMMUTATIVE COMPLETELY
PRIMARY RINGS**

Ojiema M. Onyango¹, Owino M. Oduor², Odhiambo P. Oleche³

¹Department of Mathematics
Masinde Muliro University of Science and Technology
P.O. Box 190-50100, Kakamega, KENYA

²Department of Mathematics and Computer Science
University of Kabianga
P.O. Box 2030-20200, Kericho, KENYA

³Department of Pure and Applied Mathematics
Maseno University
P.O. Box 333, Maseno, KENYA

Abstract: Let G be a group. The groups G' for which G is an automorphism group have not been fully characterized. Suppose R is a Completely Primary finite Ring with Jacobson Radical J such that $J^2 = (0)$. In this case, the characteristic of R is p or p^2 and the group of units $R^* = \mathbb{Z}_{p^r-1} \times (I + J)$. The structure of R^* is well known, but its automorphism group is not well documented. Given the group R^* , let $Aut(R^*)$ denote the group of isomorphisms $\phi : R^* \rightarrow R^*$ with multiplication given by the composition of functions. The structure of the automorphism groups of finite groups is intimately connected to the structure of the finite groups themselves. In this note, we determine the structure of $Aut(R^*)$ using well known procedures and to this end, extend the results previously obtained in this area of research.

AMS Subject Classification: 20K30, 16P10

Key Words: automorphism groups, unit groups, square radical zero, completely primary rings

1. Introduction

The definition of terms and standard notations can be obtained from [2, 3, 5, 6]. The classification of finite rings has been studied with great success in the recent past ([2, 5, 6] and related studies). Most of the researchers have concentrated in obtaining the structures of the unit groups of Completely Primary finite rings. However, the automorphisms of these unit groups have remained uncharacterized. The first general structure result for the automorphisms group of a finite group follows from a classical result of Gauss in number theory. Let \mathbb{Z}_n denote the additive group of integers mod n and $U(\mathbb{Z}_n)$ the multiplicative group of integers mod n . Gauss analyzed the orders of elements in $U(\mathbb{Z}_{p^n})$ for p prime. His results can be summarized as follows:

Theorem 1. (Gauss) *Let p be an odd prime and $n \geq 1$ or $p = 2$ and $n \geq 2$. Then*

$$U(\mathbb{Z}_{p^n}) \cong \mathbb{Z}_{p^{n-1}(p-1)}, U(\mathbb{Z}_{2^n}) \cong \mathbb{Z}_2 \times \mathbb{Z}_{2^{n-2}}.$$

Notice that $U(\mathbb{Z}_n)$ is precisely the set of generators of \mathbb{Z}_n . Since any automorphism $\theta \in \mathbb{Z}_n$ sends 1 to a generator, the valuation map $E : \text{Aut}(\mathbb{Z}_n) \mapsto U(\mathbb{Z}_{p^n})$ given by $E(\theta) = E(1)$ is an isomorphism of groups. This sets the stage for prime factorization of the integer n and consequently the classification of the automorphisms of an arbitrary finite abelian group. On the other hand, the automorphisms of cyclic groups are precisely known. In fact, given any prime p and any integer n , the group $\text{Aut}(C_p^n) \cong \text{Aut}(\mathbb{Z}_p^n)$, the group of n by n invertible matrices over the field \mathbb{Z}_p . These and related matrix groups play important roles in the classification of simple groups.

In [5], we constructed a class of Square Radical Zero Commutative Completely Primary finite Rings as follows:

Let R_o be the Galois ring of the form $GR(p^{kr}, p^k)$, such that $k = 1, 2$. For each $i = 1, \dots, h$, let $u_i \in J(R)$, such that U is an h - dimensional R_o -module generated by $\{u_1, \dots, u_h\}$ so that $R = R_o \oplus U$ is an additive group. On this group, define multiplication by the following relation

$$pu_i = u_i u_j = u_j u_i = 0, \quad u_i r_o = (r_o)^{\sigma_i} u_i, \quad (*)$$

where $r_o \in R_o$, $1 \leq i, j \leq h$, p is a prime integer, n and r are positive integers and σ_i is the automorphism associated with u_i . Further, let the generators $\{u_i\}$ for U satisfy the additional condition that, if $u_i \in U$, then, $pu_i = u_i u_j = 0$. From the given multiplication in R , we see that if $r_0 + \sum_{i=1}^h \lambda_i u_i$ and $s_0 + \sum_{i=1}^h \gamma_i u_i$,

$r_0, s_0 \in R_0, \lambda_i, \gamma_i \in F_o$ are elements of R , then,

$$(r_0 + \sum_{i=1}^h \lambda_i u_i)(s_0 + \sum_{i=1}^h \gamma_i u_i) = r_0 s_0 + \sum_{i=1}^h \{(r_0 + pR_o)\gamma_i + \lambda_i(s_0 + pR_o)^{\sigma_i}\} u_i.$$

It is easy to verify that the given multiplication turns the additive abelian group R , into a ring with identity $(1, 0, \dots, 0)$. Moreover, $J^2 = (0)$. Accordingly, the characteristic of R is either p or p^2 . Furthermore, the group of units R^* of R is given by $R^* = \mathbb{Z}_{p^{r-1}} \times (1 + J)$, a direct product of abelian groups.

In [3], Hillar and Rhea have given a useful description of the automorphism group of an arbitrary finite abelian group and they found the size of this automorphism group. We extend their work by characterizing $Aut(R^*)$. We find all the elements of $GL_{hr}(\mathbb{Z}_p)$ that can be extended to a matrix in $End(B_p)$ and calculate the distinct ways of extending such elements to the endomorphism. The first complete characterization of the automorphism group of an abelian group was however given by Ranum [1].

2. Preliminaries

Theorem 2. (cf. [5]) *The unit group R^* of the commutative completely primary finite ring of characteristic p or p^2 with maximal ideal J such that $J^2 = (0)$ and with invariants p (prime integer), $p \in J, r \geq 1$ and $h \geq 1$ is a direct product of cyclic groups as follows:*

(i) *If Char $R = p$, then*

$$R^* = \mathbb{Z}_{p^{r-1}} \times (\mathbb{Z}_p^r)^h.$$

(ii) *If Char $R = p^2$, then*

$$R^* = \mathbb{Z}_{p^{r-1}} \times \mathbb{Z}_p^r \times (\mathbb{Z}_p^r)^h.$$

The following lemma will be useful in the sequel.

Lemma 1. (cf. [3]) *Let H and K be finite groups of relatively prime orders. Then, $Aut(H) \times Aut(K) \cong Aut(H \times K)$*

Remark 1. Since the groups $\mathbb{Z}_{p^{r-1}}$ and $((\mathbb{Z}_p^r)^h)$ are of relatively prime orders and $Aut(\mathbb{Z}_{p^{r-1}}) \cong (\mathbb{Z}_{p^{r-1}})^*$, it implies that, when $\text{char}(R) = p$ then

$$Aut(R^*) \cong Aut(\mathbb{Z}_{p^{r-1}}) \times Aut((\mathbb{Z}_p^r)^h) \cong (\mathbb{Z}_{p^{r-1}})^* \times Aut((\mathbb{Z}_p^r)^h).$$

Similarly, when $\text{char}(R) = p^2$, then

$$Aut(R^*) \cong (\mathbb{Z}_{p^{r-1}})^* \times Aut((\mathbb{Z}_p^r)^{(h+1)}).$$

Lemma 2. Let $\text{Char}(R) = p$, p a prime integer and

$$B_p = 1 + J = (\mathbb{Z}_p^r)^h.$$

Then, $|B_p| = p^{rh}$.

Proof. If $G = \prod_{i=1}^r \mathbb{Z}/p^{e_i}\mathbb{Z}$ such that $1 \leq e_1 \leq e_2 \leq \dots \leq e_r$, then

$$\left| \prod_{i=1}^r \mathbb{Z}/p^{e_i}\mathbb{Z} \right| = p^{\sum_{i=1}^r e_i}.$$

Now, in B_p , $e_1 = e_2 = \dots = e_r = 1$ and there are h -tuples of such r factors of e_i , so that $(\sum_{i=1}^r e_i)h = rh$. Thus $|B_p| = p^{rh}$ as required. \square

Remark 2. Now, suppose $G \cong \prod_i^h B'_p$ such that

$$B'_p = \underbrace{\mathbb{Z}_p \times \mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r$$

over distinct set of primes p , then $\text{Aut}(G) = \prod \text{Aut}(B'_p)$.

In the sequel, we determine $\text{Aut}(I + J)$ for both the characteristics of R . Suppose $M_p = \text{End}(B_p)$, then, the elements of M_p are group homomorphisms $\alpha : B_p \rightarrow B_p$ with ring multiplication given by composition and addition taken naturally. These rings behave like matrix rings.

3. The Endomorphism Ring and its Properties

3.1. Characteristic of $R = p$

Proposition 1. Let R be a finite ring whose additive group $(R, +)$ is of type $(p^{e_1}, p^{e_2}, \dots, p^{e_l}) : e_1 \geq e_2 \geq \dots \geq e_l$. Then, R can be identified with a subring of the endomorphism ring say B of the additive group. The ring B can be considered as the ring of all $l \times l$ matrices (a_{ij}) such that $1 \leq i, j \leq l$ of the form.

$$(a_{i,j}) = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1l} \\ p^{e_1 - e_2} a_{21} & a_{22} & \cdots & a_{2l} \\ \vdots & \vdots & \vdots & \vdots \\ p^{e_1 - e_l} a_{l1} & \cdots & \cdots & a_{ll} \end{pmatrix}$$

such that

$$a_{ij} = \begin{cases} a_{ij}, & i \leq j; \\ p^{e_j - e_i} a_{ij}, & i > j. \end{cases}$$

Proposition 2. *Let*

$$1 + J = B_p = \underbrace{\mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \dots \times \mathbb{Z}_p^r}_h$$

and $B'_p = \mathbb{Z}_p^r$. We define the endomorphism ring of B'_p by

$$End(B'_p) = \{(a_{ij}) \in \mathbb{Z}_{r \times r} : p^{e_i - e_j} \mid a_{ij}; 1 \leq j \leq i \leq r\}.$$

Example 1. If $n = 4$ and since $e_1 = e_2 = e_3 = e_4 = 1$, it follows that if $a_{ij} \in End(B'_p)$ satisfies the conditions of the above proposition, that is

$$p^{e_i - e_j} \mid a_{ij}; 1 \leq j \leq i \leq r,$$

then $a_{ij} \equiv 0 \pmod{p} \forall i, j \leq 4$. Suppose $M'_p = End(B'_p)$, then

$$M'_p = \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}.$$

Thus, generally,

$$M'_p = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1r} \\ a_{21} & a_{22} & \cdots & a_{2r} \\ \vdots & \vdots & \vdots & \vdots \\ a_{r1} & a_{r2} & \cdots & a_{rr} \end{pmatrix} = End(B'_p).$$

Proposition 3. *Consider $B'_p = \underbrace{\mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r$ as defined above. Then:*

- (i) $End(B'_p) = M_r(\mathbb{Z}_p)$;
- (ii) $Aut(B'_p) = GL_r(\mathbb{Z}_p)$;
- (iii) $|Aut(B'_p)| = \prod_{i=0}^{r-1} (p^r - p^i)$.

Proof. The proof of (i) and (ii) follow from the previous proposition.

Now, consider $Aut(\underbrace{\mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r)$. We start with $Aut(\mathbb{Z}_p)$ and $Aut(\mathbb{Z}_p \times \mathbb{Z}_p)$

in order to obtain the size of the automorphism group of B'_p . In \mathbb{Z}_p , each of the $p - 1$ nonidentity elements has order p . Suppose $\mathbb{Z}_p = \langle a \rangle$, then the map $a \mapsto a^i$ is an element of $Aut(\mathbb{Z}_p)$ provided $i \in [1, p - 1]$. Thus $|Aut(\mathbb{Z}_p)| = p - 1 = \Phi(p)$, where Φ is the Euler's-phi function.

Next, let a and b each generate groups of order p , so that $\mathbb{Z}_p \times \mathbb{Z}_p = \langle a \rangle \times \langle b \rangle$. A homomorphism $\theta : \mathbb{Z}_p \times \mathbb{Z}_p \mapsto \mathbb{Z}_p \times \mathbb{Z}_p$ is an automorphism iff $|\theta(a)| = |\theta(b)| = p$ and $\langle \theta(a) \rangle$ intersects with $\langle \theta(b) \rangle$ only at identity.

To find $|\mathbb{Z}_p \times \mathbb{Z}_p|$, we must count the pairs (β, β') of elements in $\mathbb{Z}_p \times \mathbb{Z}_p$ such that $\theta(a) = \beta$ and $\theta(b) = \beta'$ determines an automorphism. Each of the $p^2 - 1$ nonidentity elements of $\mathbb{Z}_p \times \mathbb{Z}_p$ has order p , so, a given element of $Aut(\mathbb{Z}_p \times \mathbb{Z}_p)$ may map a to any of the $p^2 - 1$ different places.

Let β be nonidentity element. We must count the elements β' of $\mathbb{Z}_p \times \mathbb{Z}_p$ such that $\beta' = p$ and $\langle \beta \rangle \cap \langle \beta' \rangle = \{e\}$. Since each β generates a group of order p and any of the $p^2 - p$ elements of $\mathbb{Z}_p \times \mathbb{Z}_p$ lying outside of $\langle \beta \rangle$ will generate a group of order p that intersects the group $\langle \beta \rangle$ only at identity element, it follows that

$$|Aut(\mathbb{Z}_p \times \mathbb{Z}_p)| = (p^2 - 1)(p^2 - p).$$

For $B'_p = \underbrace{\mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r$, let $\{g_1, \dots, g_r\}$ be a set of generators for B'_p , so that

$$\underbrace{\mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r = \langle g_1 \rangle \times \langle g_2 \rangle \times \dots \times \langle g_r \rangle.$$

Each of the nonidentity elements of B'_p has order p . We now count the number of injective maps from the above generators to nonidentity elements that generate groups intersecting only at the identity element.

Suppose that an automorphism of B'_p sends g_1 to some element β in B'_p , then there are $p^r - p$ elements β' such that $\langle \beta \rangle \cap \langle \beta' \rangle = \{e\}$. Supposing further that this automorphism is given by $g_1 \mapsto \beta$ and $g_2 \mapsto \beta'$ for some β' not in $\langle \beta \rangle$, there remain $p^r - p^2$ elements $\beta'' \in B'_p$ that are outside of $\langle \beta \rangle \times \langle \beta' \rangle$. Sending g_3 to any such β'' gives $(\langle \beta \rangle \times \langle \beta' \rangle) \cap \langle \beta'' \rangle = \{e\}$.

Continuing in this manner, it is easy to specify where an automorphism of B'_p sends the first n generators and then find $p^r - p^n$ elements in B'_p to which

the next generators might be sent. Thus

$$|Aut(\prod_{i=1}^r \mathbb{Z}_p)| = |Aut(B'_p)| = \prod_{i=0}^{r-1} (p^r - p^i). \quad \square$$

Before stating our main results, we need the following properties:

From [3], we have H_p as a form of B_p thus B_p forms a ring.

Let $\theta_i : \mathbb{Z} \mapsto (\mathbb{Z}_p^r)^h$ defined by $\theta(x) = x(mod p)$ be a homomorphism given by

$$\theta(x_{11}, \dots, x_{hr}) = \theta_1(x_{11}) \dots \theta_{hr}(x_{hr}).$$

Here is the description of $End(B_p)$ as a quotient of the matrix ring M_p

Theorem 3. (See also [3]) *The map $\theta : M_p \rightarrow End(B_p)$ given by $\theta(M_p)(\alpha_{11}, \dots, \alpha_{hr})^T = \theta(M_p(\alpha_{11}, \dots, \alpha_{hr})^T)$ is a surjective ring homomorphism.*

Lemma 3. *Let K be the set of matrices $M = (a_{ij}) \in B_p$ such that $p \mid a_{ij} \forall i, j$. Then, K forms an ideal.*

The following is a complete description of $Aut(B_p)$

Lemma 4. *An endomorphism $M' = \theta(M)$ is an automorphism if and only if $M(mod p) \in GL_{hr}(\mathbb{Z}_p)$.*

3.2. Characteristic of $R = p^2$

Proposition 4. *Consider $B'_p = \underbrace{\mathbb{Z}_p \times \dots \times \mathbb{Z}_p}_r$ and $B_p = \underbrace{(\mathbb{Z}_p \times \dots \times \mathbb{Z}_p)}_r^{h+1}$.*

Then:

- (i) $End(B'_p) = M_r(\mathbb{Z}_p)$;
- (ii) $Aut(B'_p) = GL_r(\mathbb{Z}_p)$;
- (iii) $|Aut(B'_p)| = \prod_{i=0}^{r-1} (p^r - p^i)$.

Proof. Similar to the case when the characteristic of $R = p$, with some slight modifications. □

4. Main Results

Lemma 5. Consider R^* when the characteristic of $R = p$,

$$B_p = \underbrace{\mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \dots \times \mathbb{Z}_p^r}_h$$

and $M_p = \text{End}(B_p)$. Then:

(i) $\text{End}(B_p) = M_{hr}(\mathbb{Z}_p)$

$$= \begin{pmatrix} a_{11} & \cdots & a_{1r} & a_{1(r+1)} & \cdots & a_{1(2r)} & \cdots & a_{1(hr)} \\ a_{21} & \cdots & a_{2r} & a_{2(r+1)} & \cdots & a_{2(2r)} & \cdots & a_{2(hr)} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{r1} & \cdots & a_{rr} & a_{r(r+1)} & \cdots & a_{r(2r)} & \cdots & a_{r(hr)} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{(2r)1} & \cdots & a_{(2r)r} & a_{(2r)(r+1)} & \cdots & a_{(2r)(2r)} & \cdots & a_{(2r)(hr)} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{(hr)1} & \cdots & a_{(hr)r} & a_{(hr)(r+1)} & \cdots & a_{(hr)(2r)} & \cdots & a_{(hr)(hr)} \end{pmatrix};$$

(ii) $\text{Aut}(B_p) = GL_{hr}(\mathbb{Z}_p)$;

(iii) $|\text{Aut}(B_p)| = \prod_{i=0}^{(hr)-1} (p^{hr} - p^i)$.

Proof. Using our characterization and appealing to Hillar and Rhea's account in [3], we find all the elements of $GL_{hr}(\mathbb{Z}_p)$ that can be extended to a matrix in $\text{End}(B_p)$ and calculate the distinct ways of extending such an element to an endomorphism. So, we need all such matrices $M_{hr} \in \text{End}(B_p)$ that are invertible modulo p .

Now, recall that $B_p = (\mathbb{Z}_p^r)^h$ and define the following numbers:

$$d_{hr} = \max\{\eta : e_\eta = e_{hr}\}, c_{hr} = \min\{\eta : e_\eta = e_{hr}\}.$$

Since $e_\eta = e_{hr}$ for $\eta = hr$, we have the two inequalities $d_{hr} \geq hr$ and $c_{hr} \leq hr$. But in this case, $e_1 = e_2 = \dots = e_{hr}$. Therefore, $d_{hr} = hr$ and $c_{hr} = i, \forall i = 1, \dots, hr$.

Thus $M_{hr} \in \text{End}(B_p)$ are upper block triangular matrices of the form

$$\begin{pmatrix} \eta_{1d_{11}} & \cdots & \eta_{1r} & \eta_{1(r+1)} & \cdots & \eta_{1(2r)} & \cdots & \eta_{1(hr)} \\ \eta_{21} & \cdots & \eta_{2r} & \eta_{2(r+1)} & \cdots & \eta_{2(2r)} & \cdots & \eta_{2(hr)} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \eta_{rd_{11}} & \cdots & \eta_{rr} & \eta_{r(r+1)} & \cdots & \eta_{r(2r)} & \cdots & \eta_{r(hr)} \\ \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ & & & \eta_{(2r)(r+1)} & \cdots & \eta_{(2r)(2r)} & \cdots & \eta_{(2r)(d_{hr})} \\ & & & & \ddots & & & \vdots \\ 0 & & & & & & \ddots & \eta_{(hr)(d_{hr})} \end{pmatrix}.$$

The number of such M_{hr} is $\prod_{i=0}^{(hr)-1} (p^{hr} - p^i)$ since we require linearly independent columns. So, the first step for calculating $|\text{Aut}(B_p)|$ is done.

Next, to extend each entry η_{ij} from $\eta_{ij} \in (\mathbb{Z}_{p^{e_i}})^h$ to $a_{ij} \in (\mathbb{Z}_{p^{e_i}})^h$ such that $a_{ij} \equiv \eta_{ij} \pmod{p}$, since $e_1 = e_2 = \dots = e_{hr} = 1, \forall i, j$, we have p^{e_i} ways to do so for the necessary zeros (That is, when $e_i > e_j$). Similarly, there are p^{e_i-1} ways for the not necessarily zero entries, that is, when $e_i = e_j$ as any element of $(\mathbb{Z}_{p^{e_i}})^h$ works. □

Lemma 6. Consider R^* when the characteristic of $R = p^2$,

$$B_p = \underbrace{\mathbb{Z}_p^r \times \mathbb{Z}_p^r \times \dots \times \mathbb{Z}_p^r}_{h+1}$$

and $M_p = \text{End}(B_p)$. Then:

(i) $\text{End}(B_p) = M_{(h+1)r}(\mathbb{Z}_p) =$

$$\begin{pmatrix} a_{11} & \cdots & a_{1r} & a_{1(r+1)} & \cdots & a_{1(2r)} & \cdots & a_{1(h+1)r} \\ a_{21} & \cdots & a_{2r} & a_{2(r+1)} & \cdots & a_{2(2r)} & \cdots & a_{2(h+1)r} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{r1} & \cdots & a_{rr} & a_{r(r+1)} & \cdots & a_{r(2r)} & \cdots & a_{r(h+1)r} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{(2r)1} & \cdots & a_{(2r)r} & a_{(2r)(r+1)} & \cdots & a_{(2r)(2r)} & \cdots & a_{(2r)(h+1)r} \\ \vdots & \cdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{((h+1)(r))1} & \cdots & a_{((h+1)(r))r} & a_{((h+1)(r))(r+1)} & \cdots & a_{(hr)(2r)} & \cdots & a_{((h+1)r)((h+1)r)} \end{pmatrix};$$

(ii) $\text{Aut}(B_p) = GL_{(h+1)r}(\mathbb{Z}_p)$;

(iii) $|\text{Aut}(B_p)| = \prod_{i=0}^{((h+1)r)-1} (p^{(h+1)r} - p^i)$.

Proof. Follows from the proof of the previous lemma with some slight modifications. \square

Theorem 4. *Let R^* be the unit group of a class of finite rings described by the construction in Section 1. Then:*

(i) *When the characteristic of $R = p$, $\text{Aut}(R^*) = \text{Aut}(\mathbb{Z}_{p^r} - 1) \times \text{Aut}(B_p)$,*

$$|\text{Aut}(R^*)| = |\mathbb{Z}_{p^r}^* - 1| \times |\text{Aut}(B_p)| = \Phi(p^r - 1) \cdot \prod_{i=0}^{(hr)-1} (p^{hr} - p^i);$$

(ii) *When the characteristic of $R = p^2$, then*

$$|\text{Aut}(R^*)| = \Phi(p^r - 1) \cdot \prod_{i=0}^{((h+1)r)-1} (p^{(h+1)r} - p^i).$$

Acknowledgements

The first author acknowledges the co-authors for their meaningful and well done PhD supervision. Their selflessness in giving scholarly comments and direction is invaluable. May God bless you. A lot of thanks to Masinde Muliro University of Science and Technology for their financial support towards my research stay at the University of Kwazulu Natal, South Africa.

References

- [1] A. Ranum, The group of classes of congruent matrices with application to the group of isomorphisms of any Abelian groups, *Trans. Amer. Math. Soc.*, **8** (1907), 71-91, **doi:** 10.1090/S0002-9947-1907-1500775-1.
- [2] C.J. Chikunji, On unit groups of completely primary finite rings, *Mathematical Journal of Okayama University*, **50** (2008), 149-160.
- [3] C. Hillar, D. Rhea, Automorphisms of an Abelian p -group, *Amer. Math. Monthly*, **114** (2007), 917-922.
- [4] K. Shoda, Uber die Automorphismen einer endlichen Abelschen Gruppe, *Math. Ann.*, **100** (1928), 674-686, **doi:** 10.1007/BF01448871.
- [5] M.O. Oduor, M.O. Ojiema, M. Eliud, Units of commutative completely primary finite rings of characteristic p^n , *International Journ. of Algebra*, **7**, No. 6 (2013), 259-266.
- [6] Y. Alkamees, Finite Rings in which the multiplication of any two zero divisors is zero, *Arch. Math.*, **37** (1981), 144-149, **doi:** 10.1007/BF01234337.